

Emerging from the vortex

17 February 2012

A theory that accurately describes the properties of relativistic electron vortex beams will impact applications such as electron microscopy

Whether a car or a ball, the forces acting on a body moving in a straight line are very different to those acting on one moving in tight curves. This maxim also holds true at microscopic scales. As such, a beam of electrons that moves forward linearly has different properties to one with vortex-like properties. Since vortex beams show properties in magnetic fields that could lead to novel applications, a RIKEN-led research team has developed a theory that provides an understanding of these properties¹.

"Ours is the first comprehensive theory of relativistic electron vortex beams and adds significantly to their understanding," comments team member Konstantin Bliokh from the RIKEN Advanced Science Institute (ASI).

Like any particle, electrons can exhibit wave-like characteristics; and, understanding this behavior is critical to understanding the behavior of vortex beams so they can be harnessed in future applications. Unlike the broad front of an ocean wave hitting a beach, however, the oscillations of electron waves are out of sync along the beam: the slight shifts in their timing gives the waves a corkscrew character.

A vortex beam shows unique and potentially exploitable quantum effects arising from the interplay between the so-called 'orbital angular momentum' of its electrons and their intrinsic property called spin. The potential of these beams became apparent only recently, when they were demonstrated for the first time by Masaya Uchida and Akira Tonomura from ASI².

The complex mixture of electron spin, beam vortex properties and the relativistic properties of the electrons has complicated the theoretical understanding of the beams, says team leader Franco Nori also from ASI. He explains that their fundamental theoretical description was possible only through the combined consideration of quantum and relativistic properties of the electrons in the beam. This provided new insights into the interaction between the electrons' spin and the vortex property of the beam. In particular, the researchers found that this so-called spin-orbit interaction results in a different behavior for vortex beams made of electrons with spins pointing up or down, respectively—an effect that should be observable (Fig. 1).

Beyond providing these fundamental insights, the new theory also has sound practical implications, as the beams are very sensitive to magnetic fields, according to Nori. "The theoretical understanding that we have reached will eventually contribute to the development of enhanced electron microscopes that can image magnetic materials with atomic resolution," he says.

The corresponding author for this highlight is based at the [Digital Materials Team, RIKEN Advanced Science Institute](#)

1. Bliokh, K.Y., Dennis, M.R. & Nori, F. Relativistic electron vortex beams: angular momentum and spin-orbit interaction. *Physical Review Letters* **107**, 174802 (2011). [article](#)
2. Uchida, M. & Tonomura, A. Generation of electron beams carrying orbital angular momentum. *Nature* **464**, 737–739 (2010). [highlight](#)




Figure 1: A comparison of electron beams with parallel (top) and antiparallel (bottom) spin and vortex orientation. The plot shows the density (grey) and current distributions (arrows) for electrons in the x and y direction, with either 'up' (+) or 'down' (-) spin, respectively. For electrons in the center of the beam, the distribution is predicted to be different for both cases.

Reproduced, with permission, from Ref. 1 © 2012 American Physical Society