



FAST TRACK COMMUNICATION

Continuous dissipative phase transitions with or without symmetry breaking

OPEN ACCESS

RECEIVED
27 August 2021REVISED
24 November 2021ACCEPTED FOR PUBLICATION
26 November 2021PUBLISHED
22 December 2021Original content from
this work may be used
under the terms of the
Creative Commons
Attribution 4.0 licence.Any further distribution
of this work must
maintain attribution to
the author(s) and the
title of the work, journal
citation and DOI.Fabrizio Minganti^{1,2,*} , Ievgen I Arkhipov³ , Adam Miranowicz^{1,4}  and
Franco Nori^{1,5,6,*} ¹ Theoretical Quantum Physics Laboratory, RIKEN Cluster for Pioneering Research, Wako-shi, Saitama 351-0198, Japan² Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), CH-1015 Lausanne, Switzerland³ Joint Laboratory of Optics of Palacký University and Institute of Physics of CAS, Faculty of Science, Palacký University, 17. listopadu 12, 771 46 Olomouc, Czech Republic⁴ Institute of Spintronics and Quantum Information, Faculty of Physics, Adam Mickiewicz University, 61-614 Poznań, Poland⁵ RIKEN Center for Quantum Computing (RQC), Wakoshi, Saitama 351-0198, Japan⁶ Physics Department, The University of Michigan, Ann Arbor, MI 48109-1040, United States of America

* Authors to whom any correspondence should be addressed.

E-mail: fabrizio.minganti@gmail.com, ievgen.arkhipov@upol.cz, miran@amu.edu.pl and fnori@riken.jp

Keywords: phase transitions, Scully–Lamb laser, open system symmetries

Abstract

The paradigm of second-order phase transitions (PTs) induced by spontaneous symmetry breaking (SSB) in thermal and quantum systems is a pillar of modern physics that has been fruitfully applied to out-of-equilibrium open quantum systems. Dissipative phase transitions (DPTs) of second order are often connected with SSB, in close analogy with well-known thermal second-order PTs in closed quantum and classical systems. That is, a second-order DPT should disappear by preventing the occurrence of SSB. Here, we prove this statement to be wrong, showing that, surprisingly, SSB is not a necessary condition for the occurrence of second-order DPTs in *out-of-equilibrium open quantum systems*. We analytically prove this result using the Liouvillian theory of DPTs, and demonstrate this anomalous transition in a paradigmatic laser model, where we can arbitrarily remove SSB while retaining criticality, and on a Z_2 -symmetric model of a two-photon Kerr resonator. This new type of PT cannot be interpreted as a ‘semiclassical’ bifurcation, because, after the DPT, the system steady state remains unique.

1. Introduction

The similarities and differences between quantum (or thermal) phase transitions (PTs) and dissipative phase transitions (DPTs) in open quantum systems are the subject of intense research [1–8]. Criticality and critical phenomena (e.g. hysteresis [9–12] and slowing-down [13–15]) have been predicted, observed, and characterized for first-order DPTs. Central to the characterization of second-order PTs is the role of spontaneous symmetry breaking (SSB): nonanalyticity can occur when a system symmetry is ‘broken’, i.e. the emergence of several steady (or ground) states that are not invariant anymore under the action of a given symmetry group [16–18]. SSB in open systems has been discussed in, e.g. references [19–27]. The relation between criticality, symmetries, and exotic effects have also been discussed for a wide range of models [5, 28–32].

In this article, we analytically prove that *second-order DPTs in open quantum systems can occur with or without symmetry breaking*. Similarly to other examples, where the phases of dissipative systems possess features which have no analogue in closed and thermal systems [19, 33–36], this feature can be explained by the spectral properties of a Liouvillian superoperator (i.e. the generator of the dynamics of an out-of-equilibrium open quantum system) [1, 18, 37].

We demonstrate our results with two examples. First, we consider a lasing model characterized by $U(1)$ -SSB with a second-order DPT. By adding dephasing, the $U(1)$ symmetry of the model is maintained, but its phase coherence is destroyed, thus preventing SSB. Yet, a second-order DPT takes place. Second, we

consider the Z_2 symmetry breaking in parametric down-conversion. In this case, the addition of a parity dissipator enables a DPT without Z_2 -SSB.

Similar phenomena of continuous PTs without symmetry breaking (whose explanation goes beyond the Landau theory of PTs [17, 38, 39]), can be encountered in closed systems at *equilibrium* [40, 41]. These examples are characterized by a nontrivial topological structure, e.g. topological insulators [42, 43]. As such, our work prompts the question for a generalization of topological PTs in *non-equilibrium*, non-quadratic, and bosonic systems [44–46]. Other models of open quantum systems can undergo critical phenomena without a trivial SSB. For instance, in reference [47], it was shown that a continuous PT without symmetry breaking can take place. However, such a model can also be associated with the emergence of boundary time crystals [48] (signalling its profoundly dissipative nature), and meaning that a time-invariance symmetry is broken. In other words, in this model there exist multiple (oscillating) non-decaying states.

Furthermore, the presence of a DPT without SSB has been discussed for chain of S-level spins in reference [49]. The authors demonstrate that in a regime where the semiclassical solution of a $U(1)$ symmetric model would predict SSB, purely fluctuation-induced suppression of symmetry breaking can prevent states from retaining their coherence.

Our work reveals that in *any* dissipative continuous second-order PT, SSB can be arbitrarily removed. This *always* allows obtaining a unique steady state in the ‘broken symmetry’ region. As we also discuss below, such nontrivial effect can be obtained both with the use of Hamiltonian or dissipative terms, showing the nontrivial interplay between quantum and dissipative fluctuations.

2. Criticality of open quantum systems

Under the Born and Markov approximations [50], the reduced density matrix $\hat{\rho}(t)$ of an open quantum system at time t evolves according to a Lindblad master equation ($\hbar = 1$):

$$\frac{d}{dt}\hat{\rho}(t) = \mathcal{L}\hat{\rho}(t) = -i[\hat{H}, \hat{\rho}(t)] + \sum_j \mathcal{D}[\hat{L}_j]\hat{\rho}(t), \quad (1)$$

where \hat{H} is the Hamiltonian describing the coherent part of the system evolution, \mathcal{L} is the Liouvillian superoperator [37, 50–52], and $\mathcal{D}[\hat{L}_j]$ are the so-called Lindblad dissipators, whose action is

$$\mathcal{D}[\hat{L}_j]\hat{\rho}(t) = \hat{L}_j\hat{\rho}(t)\hat{L}_j^\dagger - \frac{\hat{L}_j^\dagger\hat{L}_j\hat{\rho}(t) + \hat{\rho}(t)\hat{L}_j^\dagger\hat{L}_j}{2}. \quad (2)$$

The operators \hat{L}_j are the jump operators, and they describe how the environment acts on the system inducing loss and gain of particles, energy, and information. The steady state $\hat{\rho}_{ss}$, i.e. the state which does not evolve anymore under the action of the Liouvillian ($\partial_t\hat{\rho}_{ss} = \mathcal{L}\hat{\rho}_{ss} = 0$), is central to a system characterization. We indicate the expectation values of operators at the steady state as $\langle \hat{o} \rangle_{ss} = \text{Tr}[\hat{\rho}_{ss}\hat{o}]$.

A DPT is a discontinuous change in $\hat{\rho}_{ss}$ as a function of a single parameter [1, 18]. For instance, the medium gain rate A , defined in equation (12), plays this role in the first model considered below, and:

$$\lim_{A \rightarrow A_c} \frac{\partial^2}{\partial A^2} \hat{\rho}_{ss}(A) \rightarrow \infty, \quad (3)$$

where A_c is the critical point.

This thermodynamic nonanalyticity can be witnessed in finite-size systems, as discussed in references [1, 18] and experimentally demonstrated in references [10, 15, 53]. Criticality is accompanied by the emergence of the so-called *critical slowing down*, i.e. the appearance of infinitely-long timescales in the system dynamics. Diverging timescales can be captured by the Liouvillian spectrum, defined by

$$\mathcal{L}\hat{\rho}_j = \lambda_j\hat{\rho}_j, \quad (4)$$

λ_j (the eigenvalues) representing the decay rates and oscillation frequencies, and $\hat{\rho}_j$ (the eigenmatrices) encoding the states explored along the dynamics of \mathcal{L} .

3. Dissipative phase transitions with or without spontaneous symmetry breaking

Before dealing with a specific model, let us provide a demonstration of this novel type of criticality.

The weak symmetry of a dissipative system can be described by a superoperator \mathcal{U} such that $\mathcal{U} = \hat{J} \cdot \hat{J}^\dagger$, where $\hat{J}^\dagger = \hat{J}^{-1}$ [54]. This is always the case if we assume that \mathcal{U} defines a cyclic group (such as Z_n) and

therefore $\hat{J}^\dagger \hat{J} = \hat{1}$. Note that this is one of the most common types of symmetries that one encounters in open quantum systems, characterizing, e.g. lasing (U_1), parametric downconversion (Z_2), and Z_n for a translational invariant lattice with n sites.

The fact that the Liouvillian is symmetric, i.e. $[\mathcal{L}, \mathcal{U}] = 0$ allows partitioning the space in different *symmetry sectors*, i.e. parts of the Liouvillian space which are not connected to other parts (sectors) by the system dynamics [32, 55, 56]. Accordingly, the Liouvillian reads $\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_1 + \dots$, where \mathcal{L}_k is the evolution operator of the k th symmetry sector. As such, we can relabel the eigenvalues and eigenmatrices as $\lambda_i^{(k)}$ and $\hat{\rho}_i^{(k)}$, respectively, and write

$$\mathcal{L}_k \hat{\rho}_j^{(k)} = \lambda_j^{(k)} \hat{\rho}_j^{(k)}, \quad \text{and} \quad \mathcal{U} \hat{\rho}_j^{(k)} = \hat{J} \hat{\rho}_j^{(k)} \hat{J}^\dagger = u^{(k)} \hat{\rho}_j^{(k)}. \quad (5)$$

We order the eigenvalues of each symmetry sector in such a way that $|\operatorname{Re}[\lambda_0^{(k)}]| \leq |\operatorname{Re}[\lambda_1^{(k)}]| \leq |\operatorname{Re}[\lambda_2^{(k)}]| \leq \dots$. In this regard, $\lambda_0^{(k)}$ represents the slowest-decaying process in each symmetry sector. The steady state must always be such that $\mathcal{U} \hat{\rho}_{\text{ss}} = \hat{\rho}_{\text{ss}}$ [18]. We call this the symmetry sector for $k = 0$, and therefore $\hat{\rho}_0^{(0)} \propto \hat{\rho}_{\text{ss}}$. Thus, each eigenmatrix belonging to the symmetry sector for $k = 0$, i.e. $\hat{\rho}_j^{(0)}$, is such that $\mathcal{U} \hat{\rho}_j^{(0)} = \hat{\rho}_j^{(0)}$.

These considerations on symmetries are valid independently of the presence of a DPT, but they are fundamentally related to SSB. Generally speaking, SSB occurs when a state does not have the same symmetries, across a PT, as the theory that describes it. Such a transition is characterized by an order parameter, i.e. a quantity distinguishing phases. SSB is then signalled by the fact that the order parameter is zero (due to symmetry) in one phase and in another phase it is non-zero [57]. In an open quantum system, SSB means that several steady states emerge in a region *after* a DPT took place, and each of these states is not an eigenstate of \mathcal{U} .

Consider now a Liouvillian that, in a certain region of parameter space, admits a unique eigenvalue $\lambda_0^{(0)} = 0$ and all the other eigenvalues are such that $\lambda_0^{(j)} \neq 0$ (i.e. $\hat{\rho}_{\text{ss}} \propto \hat{\rho}_0^{(0)}$, such that $\mathcal{L} \hat{\rho}_{\text{ss}} = 0$, is the only eigenmatrix which does not evolve under the Lindblad master equation). Within this formalism, a PT with SSB means that in each symmetry sector a zero eigenvalue $\lambda_0^{(k)}$ emerges, i.e.

$$\mathcal{L} \hat{\rho}_0^{(k)} \neq 0 \quad \text{if } A < A_c, \quad \text{and} \quad \mathcal{L} \hat{\rho}_0^{(k)} = 0 \quad \text{if } A > A_c, \quad (6)$$

where the critical point A_c is defined in equation (3). Indeed, equation (6) states that for $A > A_c$ it is possible to construct new steady states (i.e. well-defined density matrices which do not evolve under the action of \mathcal{L}) of the form $\hat{\rho}_{\text{ss}}^{(k)} = \hat{\rho}_0^{(0)} + c_k [\hat{\rho}_0^{(k)} + \hat{\rho}_0^{(k)\dagger}]$, and, obviously, $\mathcal{U} \hat{\rho}_{\text{ss}}^{(k)} \neq \hat{\rho}_{\text{ss}}^{(k)}$. The role of the order parameters is then played by an operator \hat{O} such that $\operatorname{Tr}[\hat{\rho}_0^{(0)} \hat{O}] = 0$, and $\operatorname{Tr}[\hat{\rho}_0^{(k)} \hat{O}] \neq 0$. Thus, equation (6) reveals a DPT with SSB.

3.1. Preventing the spontaneous-symmetry breaking

Let us now consider a new Liouvillian

$$\mathcal{L}' = \mathcal{L} + \mathcal{D}[\hat{L}]. \quad (7)$$

If \mathcal{L} is a well-defined Liouvillian, so does \mathcal{L}' , because adding a dissipator to a Liouvillian keeps \mathcal{L}' a completely positive and trace-preserving map. Note that the following property holds

$$\mathcal{D}[\hat{L}] \hat{\rho}_j^{(k)} = \hat{L} \hat{\rho}_j^{(k)} \hat{L}^\dagger - \frac{\hat{L}^\dagger \hat{L} \hat{\rho}_j^{(k)} + \hat{\rho}_j^{(k)} \hat{L}^\dagger \hat{L}}{2} \neq 0, \quad \text{if and only if } k \neq 0, \quad (8)$$

if

$$[\hat{\rho}_j^{(0)}, \hat{L}] = [\hat{\rho}_j^{(0)}, \hat{L}^\dagger] = 0, \quad \text{and} \quad [\hat{\rho}_j^{(k)}, \hat{L}] = [\hat{\rho}_j^{(k)}, \hat{L}^\dagger] \neq 0. \quad (9)$$

All $\hat{\rho}_j^{(0)}$ must remain unchanged, while for $k \neq 0$, the dynamics of \mathcal{L} and of \mathcal{L}' must differ. But since $\lambda_j^{(0)} \leq 0$, and the dynamics must be different, it can only occur that $\lambda_j^{(0)} \neq 0$.

This operator can *always* be found. Indeed, we can choose the symmetry operator \hat{J} as a jump operator because

$$\mathcal{D}[\hat{J}] \hat{\rho}_j^{(k)} = \hat{J} \hat{\rho}_j^{(k)} \hat{J}^\dagger - \frac{\hat{J}^\dagger \hat{J} \hat{\rho}_j^{(k)} + \hat{\rho}_j^{(k)} \hat{J}^\dagger \hat{J}}{2} = \mathcal{U} \hat{\rho}_j^{(k)} - \hat{\rho}_j^{(k)} = (u^{(k)} - 1) \hat{\rho}_j^{(k)} = 0, \quad \text{if and only if } k = 0, \quad (10)$$

according to equation (5), and given $\hat{J}^\dagger \hat{J} = \hat{1}$. We conclude that *one can arbitrarily remove SSB from any second-order DPT*. This does not mean that \hat{J} is the only operator which allows removing the SSB while keeping a second-order DPT. It can also be done, e.g. by a generator of a symmetry group \hat{J} , provided that \hat{J} is continuous, as we show below on the example of a $U(1)$ model.

A remark is in order. If we were to include an additional term in the Hermitian Hamiltonian, not a Lindbladian dissipator, in equation (1), the above reasoning would remain the same. Indeed, by considering $\hat{H}' = \hat{H} + \hat{L}$ (where now \hat{L} needs to be Hermitian), the commutator $[\hat{\rho}_j^{(0)}, \hat{L}] = 0$ ensures that the dynamics is again unchanged for the sector $k = 0$.

3.2. Second-order phase transitions and spontaneous-symmetry breaking in the Landau theory

PTs do not necessarily imply an SSB. For instance, a first-order PT can occur even for non-symmetric systems [38] (this is true also in out-of-equilibrium systems [13, 18, 58–60]). There are also examples of second-order PTs without SSB, such as topological PTs, including the Berezinskii–Kosterlitz–Thouless transition in two-dimensional XY model [61] or Bose gasses [62]. In this case, however, topology dictates the change in the system properties.

The phenomenological theory of PTs, developed by Landau [38], explains second-order criticality in terms of symmetries and SSB. One phase ‘possesses’ a certain symmetry (i.e. the system state is invariant under the action of the symmetry group), while, in the SSB, phase this symmetry property is no more exhibited. One assumes that there is a thermodynamic potential F (e.g. a free-energy function) that is an analytic function of an order parameter β . The presence of a symmetry constrains the form of the potential which, up to a constant shift, will read

$$F = A\beta^2 + B\frac{\beta^4}{4}. \quad (11)$$

The order parameter β will be the one which minimizes the potential F . While A and B explicitly depend on the various parameters (such as temperature, pressure, chemical potential, etc), thermodynamic considerations on the system stability imply that $B > 0$. Since $A > 0$ implies that $\beta = 0$ minimizes F , a second-order DPT occurs when A becomes negative. At this point, the order parameter β that minimizes F is no more nonzero, and therefore a continuous, but nonanalytical, change in the system parameters occurs. Since symmetry dictates that $\beta = 0$, such a second-order PT is always connected to an SSB. The Landau theory remains applicable also for the description of the continuous *quantum* PT with SSB, where one can construct the free energy functional from a given microscopic Hamiltonian [16].

Our result, instead, indicates that SSB can be arbitrarily removed from second-order DPTs: a Landau-like theory cannot correctly capture the nature of DPTs.

4. Model I: a $U(1)$ -symmetric laser model


Let us provide an example of a DPT where the SSB can be arbitrarily removed. Consider a laser-like $U(1)$ model with $\hat{H} = \omega\hat{a}^\dagger\hat{a}$ and jump operators

$$\hat{L}_1 = \frac{\hat{a}^\dagger(2A - B\hat{a}\hat{a}^\dagger)}{2\sqrt{A}}, \quad \hat{L}_2 = \sqrt{\frac{3B}{4}}\hat{a}\hat{a}^\dagger, \quad \hat{L}_3 = \sqrt{\Gamma}\hat{a}, \quad (12)$$

where \hat{a} (\hat{a}^\dagger) is the bosonic annihilation (creation) operator, \hat{L}_1 describes the laser gain, \hat{L}_2 captures the field dephasing, and \hat{L}_3 represents the particle loss. The jump operators are characterized by the rates: A for the medium gain, B for the gain saturation, Γ for the dissipation (the inverse of the photon lifetime) changing to the frame rotating at the frequency ω , we can set $\hat{H} = 0$ [32]. This model is the celebrated Scully–Lamb laser master equation in the so-called weak-gain saturation regime [63–66], valid if $A = \mathcal{O}(\Gamma)$ and $B\langle\hat{a}\hat{a}^\dagger\rangle \ll 2A$. The limits of the validity of this approximation for the system dynamics are detailed in, e.g. references [67, 68].

The model is characterized by a $U(1)$ *weak symmetry* [69, 70], which is represented by the symmetry operator $\hat{J} = \exp(i\phi\hat{a}^\dagger\hat{a})$. Indeed, the transformation $\hat{a} \rightarrow \hat{J}\hat{a}\hat{J}^\dagger \rightarrow \hat{a}e^{i\phi}$ leaves the equation of motion unchanged, but $\langle\hat{a}^\dagger\hat{a}(t)\rangle$ is *not* conserved. Thus, $\langle\hat{a}\rangle_{ss} = 0$ holds for any finite-size system. An SSB in the thermodynamic limit is, thus, signalled by $\langle\hat{a}\rangle(t \rightarrow \infty) \neq 0$, which follows from symmetry considerations [71]. Therefore, in finite-size systems, the $U(1)$ SSB is signalled by $\langle\hat{a}\rangle(t) \neq 0$ for increasingly long time, as we increase N . To better grasp the meaning of this symmetry, let us express the eigenmatrices $\hat{\rho}_j$ of the Liouvillian and the action of the symmetry operator in equation (5) in the number (Fock) basis:

$$\hat{\rho}_j^{(k)} = \sum_{m,n} c_{m,n}^{(k)} |m\rangle\langle n| \quad \text{and} \quad \mathcal{U}\hat{\rho}_j^{(k)} = \sum_{m,n} c_{m,n}^{(k)} e^{-i\phi\hat{a}^\dagger\hat{a}} |m\rangle\langle n| e^{i\phi\hat{a}^\dagger\hat{a}} = \sum_{m,n} c_{m,n}^{(k)} e^{-i\phi(m-n)} |m\rangle\langle n| = u_j^{(k)} \hat{\rho}_j^{(k)}. \quad (13)$$



We conclude that $\exp[-i\phi(m-n)]$ must be a constant and, therefore, any eigenmatrix $\hat{\rho}_j$ in equation (13) must obey

$$\hat{\rho}_j^{(k)} = \sum_m c_m^{(k)} |m\rangle \langle m-k|, \quad (14)$$

for some constant integer $k \in \mathbb{Z}$. In other words, $\hat{\rho}_j^{(k)}$ must be an operator containing elements only on one diagonal, and different symmetry sectors occupy different upper and lower diagonals.

4.1. Second-order dissipative phase transitions with the spontaneous breaking of $U(1)$ symmetry

PTs and nonanalyticity can only emerge in the thermodynamic limit. One can exploit the infinite dimension of the bosonic Hilbert space to observe a nonanalytical change in the steady state, as discussed, in, e.g. references [18, 20, 32, 72–74]. To do so, we consider a rescaling parameter N , so that the size of the Hilbert space increases, but a meaningful observable, such as the rescaled photon number, merge far from the critical point:

$$\{A, B, \Gamma\} \rightarrow \{A, B/N, \Gamma\}. \quad (15)$$

In a laser model [63], N represents an increasing number of injected three-level atoms in the lase cavity, but each with a weaker light–matter coupling. We also refer to reference [68] for a more detailed discussion of the system thermodynamic limit and on the nature of the DPT in the Scully–Lamb laser model.

To numerically simulate the results of this model, we introduce a cutoff C in the Hilbert space, i.e. we assume that $\langle m|\hat{\rho}(t)|n\rangle = 0$ if $m > C$ or $n > C$. We then verify the convergence with the cutoff, i.e. we check that by increasing C the values do not change (within a numerical precision). By decreasing the nonlinearity (i.e. increasing N), C increases.

In figure 1(a), we plot the rescaled photon number $\langle \hat{a}^\dagger \hat{a} \rangle_{ss}/N$, obtained by numerically solving $\mathcal{L}\hat{\rho}_{ss} = 0$ (solid curves), and the results of the semiclassical approximation (dashed lines) for different values of N . As one can see, the photon number is continuous, but there is an emerging ‘elbow’ signalling the occurrence of a second-order DPT.

The breaking of the $U(1)$ symmetry means the retaining of coherence for infinitely long time. Although this occurs only in the thermodynamic limit, we can show the presence of the critical slowing down for finite-size systems. In figures 1(b)–(e), we show the Wigner function $W(\alpha)$ for an initially coherent state when $\eta = 0$ in the ‘broken symmetry region’ (i.e. for $A > A_c = \Gamma$), where $W(\alpha) = 2 \text{Tr} [\hat{D}_\alpha \exp(i\pi\hat{a}^\dagger \hat{a}) \hat{D}_\alpha^\dagger \hat{\rho}(t)] / \pi$ and $\hat{D}_\alpha = \exp(\alpha\hat{a}^\dagger - \alpha^*\hat{a})$ is the displacement operator [75]. As time passes, the system retains its coherence $\langle \hat{a}(t) \rangle \neq 0$, signalling that, even for this finite-size system, a critical timescale associated with SSB has emerged.

To better quantify the meaning of SSB and the role of the $U(1)$ symmetry, we note that

$$\text{Tr} \left[\hat{a}^n \hat{\rho}_j^{(k)} \right] = \text{Tr} \left[\mathcal{U}^\dagger \mathcal{U} \hat{a}^n \hat{\rho}_j^{(k)} \right] = \text{Tr} \left[\left(e^{+i\phi \hat{a}^\dagger \hat{a}} \hat{a}^n e^{-i\phi \hat{a}^\dagger \hat{a}} \right) \left(e^{-i\phi \hat{a}^\dagger \hat{a}} \hat{\rho}_j^{(k)} e^{i\phi \hat{a}^\dagger \hat{a}} \right) \right] = e^{-i\phi(k-n)} \left\langle \hat{a}^n \hat{\rho}_j^{(k)} \right\rangle. \quad (16)$$

We, thus, conclude that the critical slowing down, associated with SSB in the k th symmetry sector, is witnessed by $\langle \hat{a}^k \rangle$. Thus, in figures 1(i)–(k) we plot with red solid curves the dynamics of $\langle \hat{a}^\dagger \hat{a}(t) \rangle$, $|\langle \hat{a}(t) \rangle|$, and $|\langle \hat{a}^2(t) \rangle|$ (associated with the sectors $k = 0, \pm 1, \pm 2$), respectively. The initial state is the same coherent state as in figure 1(b). While the photon number rapidly converges to its steady-state value, we see again that the coherences are preserved for very long-times, confirming that indeed a $U(1)$ SSB is taking place, and there is a critical slowing down in each symmetry sector.

4.2. Removing the spontaneous breaking of $U(1)$ symmetry

According to our proof, we should be able to remove the SSB while retaining criticality. To do that, we notice that a jump operator of the form $\hat{L} = \sqrt{\eta/4} \hat{a} \hat{a}^\dagger$, where η represents an additional dephasing rate, satisfies equation (9) given the structure of $\hat{\rho}_j^{(k)}$ (cf equation (14)).

First, we verified that the photon number is identical to the one for which $\eta = 0$ (the results are within a floating-point precision in figure 1(a)). Again, the photon number becomes sharper and sharper with increasing N , and the results coincide for all tested values of η , meaning that a second-order DPT is taking place.

Figures 1(f)–(h) show that an initially coherent state, as in the case $\eta \neq 0$, rapidly loses its coherence on a timescale of $\eta/2$. That is, the $U(1)$ SSB does not take place. Indeed, $\hat{\rho}(t) \simeq \hat{\rho}_{ss}$ in figure 1(h), indicating that an initial state rapidly reaches its steady state, proving the absence of any residual or hidden SSB, which would anyhow lead to a critical slowing down.

To better quantify how each symmetry sector is affected, we compare $\langle \hat{a}^\dagger \hat{a}(t) \rangle$, $|\langle \hat{a}(t) \rangle|$, and $|\langle \hat{a}^2(t) \rangle|$ in figures 1(i)–(k) for $\eta = 0$ (red solid curves) and $\eta \neq 0$ (blue dashed curves). While the sector for $k = 0$ is unaffected (as demonstrated by the photon number). Dynamics in other symmetry sectors is much faster, and $|\langle \hat{a}^n(t) \rangle|$ rapidly reaches zero.

We have, thus, demonstrated the existence of a second-order DPT without the $U(1)$ SSB and the possibility to arbitrary remove the SSB retaining a critical behavior. Importantly, the semiclassical analysis [71] fails to capture this kind of criticality (see also reference [68] for details), which highlights the necessity to use the Liouvillian formalism in describing dissipative critical phenomena.

5. Model II: a Z_2 -symmetric Kerr resonator

To further illustrate the validity of our results, here we consider the second-order DPT of a two-photon Kerr resonator, studied in, e.g. [18, 20, 22, 28]. The Hamiltonian in the frame rotating at the pump frequency reads

$$\hat{H} = -\Delta \hat{a}^\dagger \hat{a} + i \frac{G}{2} \left[(\hat{a}^\dagger)^2 - \hat{a}^2 \right] + \frac{U}{2} (\hat{a}^\dagger)^2 \hat{a}^2, \quad (17)$$

where Δ is the cavity-to-pump detuning, G is the two-photon drive intensity, and U is the Kerr nonlinear interaction.

Photons continuously escape the Kerr resonator, and the system is described by the Lindblad master equation for the system density matrix $\hat{\rho}(t)$ [76]

$$\frac{\partial}{\partial t} \hat{\rho}(t) = -i[\hat{H}, \hat{\rho}(t)] + \Gamma \mathcal{D}[\hat{a}] \hat{\rho}(t). \quad (18)$$


This system is characterized by a Z_2 weak symmetry, meaning that $\hat{a} \rightarrow \hat{J} \hat{a} \hat{J}^\dagger = -\hat{a}$ leaves the equation of motion unchanged, where $\hat{J} = \exp(i\pi \hat{a}^\dagger \hat{a})$, but parity is not a conserved quantity [70]. With a reasoning similar to that of equation (13), we can demonstrate that there are two symmetry sectors, $k = 0$ and $k = 1$, such that

$$\mathcal{U} \hat{\rho}_j^{(k)} = e^{ik\pi} \hat{\rho}_j^{(k)}, \quad \text{and} \quad \hat{\rho}_j^{(k)} = \sum_{m,n} c_{2m,2n}^{(k)} |2m\rangle \langle 2n+k| + c_{2m+1,2n+1}^{(k)} |2m+1\rangle \langle 2m+1+k|. \quad (19)$$

We conclude that $\hat{\rho}_j^{(0)}$ contains only the even–even and odd–odd states, while $\hat{\rho}_j^{(1)}$ couples the even–odd and odd–even states.

This time, the rescaling parameter N acts as $\{\Delta, U, G, \Gamma\} \rightarrow \{\Delta, U/N, G, \Gamma\}$ [20, 77, 78].

Similarly to the previous case, we observe the emergence of a second-order DPT (figure 2(a)). This is accompanied by a critical slowing down signaling an SSB, as it can be argued from the evolution of a system



initialized in a coherent state. In particular, both the Wigner functions in figures 2(b)–(e) and the evolution of $\langle \hat{a}^\dagger \hat{a} \rangle$ and $|\langle \hat{a} \rangle|$ (solid lines in figures 2(i) and (j)) show the presence of a slow-time scale in the $k = 1$ symmetry sector.

To remove the SSB and keep the second-order DPT, this time we consider an additional jump operator of the form $\hat{L} = \sqrt{\zeta} \hat{J} = \sqrt{\zeta} \exp(i\pi \hat{a}^\dagger \hat{a})$. Again, there is no difference with the case for $\zeta = 0$ in the symmetry sector for $k = 0$, as it can be argued by the fact that the photon number in the steady state is unchanged (figure 2(a)) as well as the time dynamics of $\langle \hat{a}^\dagger \hat{a} \rangle$ (figure 2(i)). However, the presence of ζ significantly changes the sector $k = 1$, as it can be seen by analyzing the Wigner function in figures 2(f)–(h) and the time evolution of $|\langle \hat{a} \rangle|$ in figure 2(j).

We confirm again our predictions, and we show that, by adding an appropriate dissipator, we can remove the SSB also in the case of this Z_2 SSB.

6. Conclusions

In this article, we have proved that continuous DPTs can occur with and without an SSB. In particular, we derived analytical conditions to remove SSB from any second-order DPT. As examples, we have analyzed a paradigmatic non-equilibrium lasing system and a model characterized by a discrete Z_2 symmetry. In both cases, our analytical predictions are confirmed by the numerical simulations, demonstrating how tremendously multiform and various are the non-equilibrium states, their dynamics, and their PTs.

The presented DPTs without SSB are, to our knowledge, a novel phenomenon, where dissipation plays a fundamental role. This phenomenon opens questions concerning the mechanism of criticality in open quantum systems. Our predictions can be experimentally tested with, e.g. superconducting circuits, where physically engineered dissipation can be realized with state-of-the-art techniques.

From a fundamental point of view, the presence of second-order DPTs, where degeneracy can be removed, is intriguing because it represents a shift from the Landau theory of PTs. An interesting question would now be if it is possible, with similar mechanisms, to ‘reintroduce’ SSB in second-order DPTs without SSB, such as that in reference [49]. Revealing a link, if any, with extensions of topological theories for DPTs is one of our future objectives [45, 46]. Even though the models considered here were characterized by a single order parameter, our results are valid, in general, for any multi-order parameter systems with any global symmetry, e.g. open and/or dissipative extensions of thermal Z_2 -symmetric systems with light–matter interactions [79, 80].

Acknowledgments

The authors are grateful to the RIKEN Advanced Center for Computing and Communication (ACCC) for the allocation of computational resources of the RIKEN supercomputer system (HOKUSAI BigWaterfall). IA thanks the Grant Agency of the Czech Republic (Project No. 18-08874S), and Project No. CZ.02.1.010.00.016_0190000754 of the Ministry of Education, Youth and Sports of the Czech Republic. AM is supported by the Polish National Science Centre (NCN) under the Maestro Grant No. DEC-2019/34/A/ST2/00081. FN is supported in part by: Nippon Telegraph and Telephone Corporation (NTT) Research, the Japan Science and Technology Agency (JST) [via the Quantum Leap Flagship Program (Q-LEAP) program, the Moonshot R & D Grant No. JPMJMS2061, and the Centers of Research Excellence in Science and Technology (CREST) Grant No. JPMJCR1676], the Japan Society for the Promotion of Science (JSPS) [via the Grants-in-Aid for Scientific Research (KAKENHI) Grant No. JP20H00134 and the JSPS CRFBR Grant No. JPJSBP120194828], the Army Research Office (ARO) (Grant No. W911NF-18-1-0358), the Asian Office of Aerospace Research and Development (AOARD) (via Grant No. FA2386-20-1-4069), and the Foundational Questions Institute Fund (FQXi) via Grant No. FQXi-IAF19-06.

Data availability statement

All data that support the findings of this study are included within the article (and any supplementary files).

ORCID iDs

Fabrizio Minganti  <https://orcid.org/0000-0003-4850-1130>

Ievgen I Arkhipov  <https://orcid.org/0000-0001-6547-8855>

Adam Miranowicz  <https://orcid.org/0000-0002-8222-9268>

Franco Nori  <https://orcid.org/0000-0003-3682-7432>

References

- [1] Kessler E M, Giedke G, Imamoglu A, Yelin S F, Lukin M D and Cirac J I 2012 Dissipative phase transition in a central spin system *Phys. Rev. A* **86** 012116
- [2] Jing H, Özdemir Ş K, Lü X-Y, Zhang J, Yang L and Nori F 2014 \mathcal{PT} -Symmetric phonon laser *Phys. Rev. Lett.* **113** 053604
- [3] Marino J and Diehl S 2016 Driven Markovian quantum criticality *Phys. Rev. Lett.* **116** 070407
- [4] Rota R, Minganti F, Ciuti C and Savona V 2019 Quantum critical regime in a quadratically driven nonlinear photonic lattice *Phys. Rev. Lett.* **122** 110405
- [5] Lieu S, Belyansky R, Young J T, Lundgren R, Albert V V and Gorshkov A V 2020 Symmetry breaking and error correction in open quantum systems *Phys. Rev. Lett.* **125** 240405
- [6] Soriente M, Heugel T L, Arimitsu K, Chitra R and Zilberberg O 2021 Distinctive class of dissipation-induced phase transitions and their universal characteristics *Phys. Rev. Res.* **3** 023100
- [7] Roy A, Jahani S, Langrock C, Fejer M and Marandi A 2021 Spectral phase transitions in optical parametric oscillators *Nat. Commun.* **12** 835
- [8] Rossini D and Vicari E 2021 Coherent and dissipative dynamics at quantum phase transitions *Phys. Rep.* **936** 1
- [9] Casteels W, Storme F, Le Boité A and Ciuti C 2016 Power laws in the dynamic hysteresis of quantum nonlinear photonic resonators *Phys. Rev. A* **93** 033824
- [10] Rodriguez S R K *et al* 2017 Probing a dissipative phase transition via dynamical optical hysteresis *Phys. Rev. Lett.* **118** 247402
- [11] Landa H, Schiró M and Misguich G 2020 Multistability of driven-dissipative quantum spins *Phys. Rev. Lett.* **124** 043601
- [12] Landa H, Schiró M and Misguich G 2020 Correlation-induced steady states and limit cycles in driven dissipative quantum systems *Phys. Rev. B* **102** 064301
- [13] Casteels W, Fazio R and Ciuti C 2017 Critical dynamical properties of a first-order dissipative phase transition *Phys. Rev. A* **95** 012128
- [14] Fink J M, Dombi A, Vukics A, Wallraff A and Domokos P 2017 Observation of the photon-blockade breakdown phase transition *Phys. Rev. X* **7** 011012
- [15] Fink T, Schade A, Höfling S, Schneider C and Imamoglu A 2018 Signatures of a dissipative phase transition in photon correlation measurements *Nat. Phys.* **14** 365
- [16] Sachdev S 2001 *Quantum Phase Transitions* (Cambridge: Cambridge University Press)
- [17] Landau L 1936 The theory of phase transitions *Nature* **138** 840
- [18] Minganti F, Biella A, Bartolo N and Ciuti C 2018 Spectral theory of Liouvillians for dissipative phase transitions *Phys. Rev. A* **98** 042118
- [19] Lee T E, Gopalakrishnan S and Lukin M D 2013 Unconventional magnetism via optical pumping of interacting spin systems *Phys. Rev. Lett.* **110** 257204
- [20] Bartolo N, Minganti F, Casteels W and Ciuti C 2016 Exact steady state of a Kerr resonator with one- and two-photon driving and dissipation: controllable Wigner-function multimodality and dissipative phase transitions *Phys. Rev. A* **94** 033841
- [21] Kepesidis K V, Milburn T J, Huber J, Makris K G, Rotter S and Rabl P 2016 \mathcal{PT} -symmetry breaking in the steady state of microscopic gain-loss systems *New J. Phys.* **18** 095003
- [22] Savona V 2017 Spontaneous symmetry breaking in a quadratically driven nonlinear photonic lattice *Phys. Rev. A* **96** 033826

- [23] Rota R, Storme F, Bartolo N, Fazio R and Ciuti C 2017 Critical behavior of dissipative two-dimensional spin lattices *Phys. Rev. B* **95** 134431
- [24] Biella A, Storme F, Lebreuilly J, Rossini D, Fazio R, Carusotto I and Ciuti C 2017 Phase diagram of incoherently driven strongly correlated photonic lattices *Phys. Rev. A* **96** 023839
- [25] Huber J and Rabl P 2019 Active energy transport and the role of symmetry breaking in microscopic power grids *Phys. Rev. A* **100** 012129
- [26] Muñoz C S and Jaksch D 2021 Squeezed lasing *Phys. Rev. Lett.* **127** 183603
- [27] Takemura N, Takiguchi M and Notomi M 2021 Low- and high- β lasers in the class-A limit: photon statistics, linewidth, and the laser-phase transition analogy *J. Opt. Soc. Am. B* **38** 699
- [28] Bartolo N, Minganti F, Lolli J and Ciuti C 2017 Homodyne versus photon-counting quantum trajectories for dissipative Kerr resonators with two-photon driving *Eur. Phys. J. Spec. Top.* **226** 2705
- [29] Rota R, Minganti F, Biella A and Ciuti C 2018 Dynamical properties of dissipative XYZ Heisenberg lattices *New J. Phys.* **20** 045003
- [30] Sánchez Muñoz C, Buča B, Tindall J, González-Tudela A, Jaksch D and Porras D 2019 Symmetries and conservation laws in quantum trajectories: dissipative freezing *Phys. Rev. A* **100** 042113
- [31] Tindall J, Muñoz C S, Buča B and Jaksch D 2020 Quantum synchronisation enabled by dynamical symmetries and dissipation *New J. Phys.* **22** 013026
- [32] Minganti F, Arkhipov I I, Miranowicz A and Nori F 2020 Correspondence between dissipative phase transitions of light and time crystals (arXiv:2008.08075)
- [33] Jin J, Rossini D, Fazio R, Leib M and Hartmann M J 2013 Photon solid phases in driven arrays of nonlinearly coupled cavities *Phys. Rev. Lett.* **110** 163605
- [34] Jin J, Biella A, Viyuela O, Mazza L, Keeling J, Fazio R and Rossini D 2016 Cluster mean-field approach to the steady-state phase diagram of dissipative spin systems *Phys. Rev. X* **6** 031011
- [35] Diehl S, Micheli A, Kantian A, Kraus B, Büchler H P and Zoller P 2008 Quantum states and phases in driven open quantum systems with cold atoms *Nat. Phys.* **4** 878
- [36] Verstraete F, Wolf M M and Ignacio Cirac J 2009 Quantum computation and quantum-state engineering driven by dissipation *Nat. Phys.* **5** 633
- [37] Minganti F, Miranowicz A, Chhajlany R W and Nori F 2019 Quantum exceptional points of non-Hermitian Hamiltonians and Liouvillians: the effects of quantum jumps *Phys. Rev. A* **100** 062131
- [38] Landau L D and Lifshitz E M 2013 Statistical physics *Course of Theoretical Physics* vol 5 (Amsterdam: Elsevier)
- [39] Haar D T (ed) 1965 *Collected Papers of L.D. Landau* (Oxford: Pergamon)
- [40] Kitaev A, Lebedev V and Feigelman M 2009 Periodic table for topological insulators and superconductors *AIP Conf. Proc.* **1134** 22
- [41] Chen X, Gu Z-C, Liu Z-X and Wen X-G 2013 Symmetry protected topological orders and the group cohomology of their symmetry group *Phys. Rev. B* **87** 155114
- [42] Hasan M Z and Kane C L 2010 Colloquium: topological insulators *Rev. Mod. Phys.* **82** 3045
- [43] Qi X-L and Zhang S-C 2011 Topological insulators and superconductors *Rev. Mod. Phys.* **83** 1057
- [44] Bardyn C-E, Baranov M A, Kraus C V, Rico E, Imamoğlu A, Zoller P and Diehl S 2013 Topology by dissipation *New J. Phys.* **15** 085001
- [45] Gong Z, Ashida Y, Kawabata K, Takasan K, Higashikawa S and Ueda M 2018 Topological phases of non-Hermitian systems *Phys. Rev. X* **8** 031079
- [46] Lieu S, McGinley M and Cooper N R 2020 Tenfold way for quadratic Lindbladians *Phys. Rev. Lett.* **124** 040401
- [47] Hannukainen J and Larson J 2018 Dissipation-driven quantum phase transitions and symmetry breaking *Phys. Rev. A* **98** 042113
- [48] Iemini F, Russomanno A, Keeling J, Schirò M, Dalmonte M and Fazio R 2018 Boundary time crystals *Phys. Rev. Lett.* **121** 035301
- [49] Huber J, Kirtton P and Rabl P 2020 Nonequilibrium magnetic phases in spin lattices with gain and loss *Phys. Rev. A* **102** 012219
- [50] Breuer H and Petruccione F 2007 *The Theory of Open Quantum Systems* (Oxford: Oxford University Press)
- [51] Carmichael H 2007 *Statistical Methods in Quantum Optics 2 (Non-Classical Fields)* (Berlin: Springer)
- [52] Lidar D A 2019 Lecture notes on the theory of open quantum systems (arXiv:1902.00967)
- [53] Fitzpatrick M, Sundaresan N M, Li A C Y, Koch J and Houck A A 2017 Observation of a dissipative phase transition in a one-dimensional circuit QED lattice *Phys. Rev. X* **7** 011016
- [54] Baumgartner B and Narnhofer H 2008 Analysis of quantum semigroups with GKS-Lindblad generators: II. General *J. Phys. A: Math. Theor.* **41** 395303
- [55] Moodie R I, Ballantine K E and Keeling J 2018 Generalized classes of continuous symmetries in two-mode Dicke models *Phys. Rev. A* **97** 033802
- [56] Palacino R and Keeling J 2021 Atom-only theories for $U(1)$ symmetric cavity-QED models *Phys. Rev. Res.* **3** 032016
- [57] Beekman A J, Rademaker L and van Wezel J 2019 An introduction to spontaneous symmetry breaking *SciPost Phys. Lect. Notes* **11** 1-40
- [58] Vicentini F, Minganti F, Rota R, Orso G and Ciuti C 2018 Critical slowing down in driven-dissipative Bose-Hubbard lattices *Phys. Rev. A* **97** 013853
- [59] Foss-Feig M, Niroula P, Young J T, Hafezi M, Gorshkov A V, Wilson R M and Maghrebi M F 2017 Emergent equilibrium in many-body optical bistability *Phys. Rev. A* **95** 043826
- [60] Biondi M, Blatter G, Türeci H E and Schmidt S 2017 Nonequilibrium gas-liquid transition in the driven-dissipative photonic lattice *Phys. Rev. A* **96** 043809
- [61] Kosterlitz J M and Thouless D J 1973 Ordering, metastability and phase transitions in two-dimensional systems *J. Phys. C: Solid State Phys.* **6** 1181
- [62] Hadzibabic Z, Krüger P, Cheneau M, Battelier B and Dalibard J 2006 Berezinskii-Kosterlitz-Thouless crossover in a trapped atomic gas *Nature* **441** 1118
- [63] Yamamoto Y and Imamoğlu A 1999 *Mesoscopic Quantum Optics* (New York: Wiley)
- [64] Sargent M III, Scully M and Lamb W E 1974 *Laser Physics* (Boulder, CO: Westview Press)
- [65] Gea-Banacloche J 1997 Emergence of classical radiation fields through decoherence in the Scully-Lamb laser model *Found. Phys.* **28** 531
- [66] Arkhipov I I, Miranowicz A, Di Stefano O, Stassi R, Savasta S, Nori F and Özdemir Ş K 2019 Scully-Lamb quantum laser model for parity-time-symmetric whispering-gallery microcavities: gain saturation effects and nonreciprocity *Phys. Rev. A* **99** 042309
- [67] Wang Y K and Lamb W E 1973 Quantum theory of an optical maser: VI. Transient behavior *Phys. Rev. A* **8** 866

- [68] Minganti F, Arkhipov I I, Miranowicz A and Nori F 2021 Liouvillian spectral collapse in the Scully–Lamb laser model *Phys. Rev. Res.* (in press)
- [69] Buča B and Prosen T 2012 A note on symmetry reductions of the Lindblad equation: transport in constrained open spin chains *New J. Phys.* **14** 073007
- [70] Albert V V and Jiang L 2014 Symmetries and conserved quantities in Lindblad master equations *Phys. Rev. A* **89** 022118
- [71] DeGiorgio V and Scully M O 1970 Analogy between the laser threshold region and a second-order phase transition *Phys. Rev. A* **2** 1170
- [72] Carmichael H J 2015 Breakdown of photon blockade: a dissipative quantum phase transition in zero dimensions *Phys. Rev. X* **5** 031028
- [73] Casteels W and Ciuti C 2017 Quantum entanglement in the spatial-symmetry-breaking phase transition of a driven-dissipative Bose–Hubbard dimer *Phys. Rev. A* **95** 013812
- [74] Curtis J B, Boettcher I, Young J T, Maghrebi M F, Carmichael H, Gorshkov A V and Foss-Feig M 2021 Critical theory for the breakdown of photon blockade *Phys. Rev. Res.* **3** 023062
- [75] Haroche S and Raimond J M 2006 *Exploring the Quantum (Atoms, Cavities, and Photons)* (Oxford: Oxford University Press)
- [76] Wiseman H and Milburn G 2010 *Quantum Measurement and Control* (Cambridge: Cambridge University Press)
- [77] Minganti F, Bartolo N, Lolli J, Casteels W and Ciuti C 2016 Exact results for Schrödinger cats in driven-dissipative systems and their feedback control *Sci. Rep.* **6** 26987
- [78] Candia R D, Minganti F, Petrovnik K V, Paraoanu G S and Felicetti S 2021 Critical parametric quantum sensing (arXiv:2107.04503[quant-ph])
- [79] Schiró M, Joshi C, Bordyuh M, Fazio R, Keeling J and Türeci H E 2016 Exotic attractors of the nonequilibrium Rabi–Hubbard model *Phys. Rev. Lett.* **116** 143603
- [80] Bazhenov A Y, Tsarev D V and Alodjants A P 2021 Mean-field theory of superradiant phase transition in complex networks *Phys. Rev. E* **103** 062309